

10/706,659
L/cook 1/8/06
Search.

d his

(FILE 'HOME' ENTERED AT 15:54:04 ON 08 JAN 2006)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, JAPIO' ENTERED AT 15:54:22 ON 08
JAN 2006

L1 17 S IK17?
L2 8 DUPLICATE REMOVE L1 (9 DUPLICATES REMOVED)

=>

d his

(FILE 'HOME' ENTERED AT 15:54:04 ON 08 JAN 2006)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, JAPIO' ENTERED AT 15:54:22 ON 08
JAN 2006

L1 17 S IK17?

L2 8 DUPLICATE REMOVE L1 (9 DUPLICATES REMOVED)

=>

ordered
1/8/06 LVCook

ANSWER 5 OF 8 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN

DUPLICATE 3

AN 1997:112650 BIOSIS

DN PREV199799411853

TI DELTA-**IK17** antigen: A possible early marker of cancer development.

AU Perez, S. [Reprint author]; Sotiriadou, R.; Missitzis, G.; Trangas, T.; Giannoulis, N.; Mavroudeas, D.; Apostolikas, N.; Kokkinopoulos, D.

CS Dep. Immunol., Hellenic Anticancer Inst., 171 Alexandras Ave., 11522 Athens, Greece

SO International Journal of Biological Markers, (1996) Vol. 11, No. 4, pp. 190-197.

CODEN: IBMAEP. ISSN: 0393-6155.

DT Article

LA English

ED Entered STN: 10 Mar 1997

Last Updated on STN: 10 Mar 1997

AB DELTA-**IK17** is a 44 kD molecule located on the surface of T, B and NK cells in normal peripheral blood mononuclear cells (PBMC) (1). The portion of PBMC expressing DELTA-**IK17** was determined in 52 patients with benign breast diseases, 182 patients with breast malignancies and 132 healthy individuals. The percentage of DELTA-**IK17**-positive cells was significantly lower in the early stages (I-IIA) of malignancy compared to that of healthy donors. However, the percentage of PBMC expressing DELTA-**IK17** tended to increase as the stage of the disease advanced. DELTA-**IK17** seems to be the only antigen among the other cellular markers tested (CD2, CD4, CD8, HLA-DR) with a statistically significant correlation between a low percentage of positive cells and the early stages of malignancy and between a high expression and advanced disease. Its potential use as a tumor marker in breast cancer is discussed here.

CC Biochemistry studies - Proteins, peptides and amino acids 10064

Pathology - Diagnostic 12504

Neoplasms - Diagnostic methods 24001

IT Major Concepts

Oncology (Human Medicine, Medical Sciences)

IT Miscellaneous Descriptors

BIOCHEMISTRY AND BIOPHYSICS; CANCER; CANCER DEVELOPMENT EARLY MARKER;

DELTA-**IK17** ANTIGEN; NEOPLASTIC DISEASE; ONCOLOGY; PATIENT

ORGN Classifier

Hominidae 86215

Super Taxa

Primates; Mammalia; Vertebrata; Chordata; Animalia

Organism Name

human

Taxa Notes

Animals, Chordates, Humans, Mammals, Primates, Vertebrates

ANSWER 5 OF 8 BIOSIS COPYRIGHT (c) 2006 The Thomson Corporation on STN
 DUPLICATE 3
 AN 1997:112650 BIOSIS
 DN PREV199799411853
 TI DELTA-**IK17** antigen: A possible early marker of cancer
 development.
 AU Perez, S. [Reprint author]; Sotiriadou, R.; Missitzis, G.; Trangas, T.;
 Giannoulis, N.; Mavroudeas, D.; Apostolikas, N.; Kokkinopoulos, D.
 CS Dep. Immunol., Hellenic Anticancer Inst., 171 Alexandras Ave., 11522
 Athens, Greece
 SO International Journal of Biological Markers, (1996) Vol. 11, No. 4, pp.
 190-197.
 CODEN: IBMAEP. ISSN: 0393-6155.
 DT Article
 LA English
 ED Entered STN: 10 Mar 1997
 Last Updated on STN: 10 Mar 1997
 AB DELTA-**IK17** is a 44 kD molecule located on the surface of T, B
 and NK cells in normal peripheral blood mononuclear cells (PBMC) (1). The
 portion of PBMC expressing DELTA-**IK17** was determined in 52
 patients with benign breast diseases, 182 patients with breast
 malignancies and 132 healthy individuals. The percentage of DELTA-
IK17-positive cells was significantly lower in the early stages
 (I-IIA) of malignancy compared to that of healthy donors. However, the
 percentage of PBMC expressing DELTA-**IK17** tended to increase as
 the stage of the disease advanced. DELTA-**IK17** seems to be the
 only antigen among the other cellular markers tested (CD2, CD4, CD8,
 HLA-DR) with a statistically significant correlation between a low
 percentage of positive cells and the early stages of malignancy and
 between a high expression and advanced disease. Its potential use as a
 tumor marker in breast cancer is discussed here.
 CC Biochemistry studies - Proteins, peptides and amino acids 10064
 Pathology - Diagnostic 12504
 Neoplasms - Diagnostic methods 24001
 IT Major Concepts
 Oncology (Human Medicine, Medical Sciences)
 IT Miscellaneous Descriptors
 BIOCHEMISTRY AND BIOPHYSICS; CANCER; CANCER DEVELOPMENT EARLY MARKER;
 DELTA-**IK17** ANTIGEN; NEOPLASTIC DISEASE; ONCOLOGY; PATIENT
 ORGN Classifier
 Hominidae 86215
 Super Taxa
 Primates; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 human
 Taxa Notes
 Animals, Chordates, Humans, Mammals, Primates, Vertebrates